

- King, E. J. (1932). The colorimetric determination of phosphorus. *Biochemical Journal*, 26(2), 292-297.
- Lacava, P. T., Machado, P. C., & de Andrade, P. H. M. (2021). Phosphate Solubilization by Endophytes from the Tropical Plants. *Endophytes: Mineral Nutrient Management, Volume 3*, 207-226.
- Lale, G., & Gadre, R. (2010). Enhanced production of gibberellin A4 (GA4) by a mutant of *Gibberella fujikuroi* in wheat gluten medium. *Journal of Industrial Microbiology and Biotechnology*, 37(3), 297-306.
- Mondal, S., Halder, S. K., Yadav, A. N., and Mondal, K. C. (2020). Microbial Consortium with Multifunctional Plant Growth-Promoting Attributes: Future Perspective in Agriculture. *Advances in plant microbiome and sustainable agriculture*, 219-258.
- Mushtaq, Z., Faizan, S., and Hussain, A. (2021). Role of Microorganisms as Biofertilizers. In *Microbiota and Biofertilizers* 83-98. Springer, Cham.
- Pandya, N. D., and Desai, P. V. (2013). Gibberellic Acid Production by *Bacillus cereus* Isolated from the Rhizosphere of Sugarcane. *Journal of Pure and Applied Microbiology*, 7(4), 3239-3242.
- Patel, T., & Saraf, M. (2017). Biosynthesis of phytohormones from novel rhizobacterial isolates and their in vitro plant growth-promoting efficacy. *Journal of Plant Interactions*, 12(1), 480-487.
- Pathak, R., Paudel, V., Shrestha, A., Lamichhane, J., and Gauchan, D. P. (2017). Isolation of Phosphate Solubilizing Bacteria and their Use for Plant Growth Promotion in Tomato Seedling and Plant. *Kathmandu University Journal of Science, Engineering and Technology*, 13(2), 61-70.
- [Sagervanshi A.](#), [Kumari P.](#), [Nagee A.](#) and [Kumar A.](#) (2012). Isolation and characterization of phosphate solubilizing bacteria from Anand agriculture soil. *Advance Research in Pharmaceuticals and Biologicals*, 2(3).
- Salwan, R., Sharma, V., Saini, R., & Pandey, M. (2021). Identification of plant beneficial *Bacillus* spp. for Resilient agricultural ecosystem. *Current Research in Microbial Sciences*, 2, 100046.
- Sangeetha, V., and Thevanathan, R. (2010). Biofertilizer potential of traditional and panchagavya amended with seaweed extract. *The Journal of American Science*, 6 (2), 61-67.
- Seshagiri S., & Tallapragada, P. (2016). Study of acid phosphatase in solubilization of inorganic phosphates by *Piriformospora indica*. *Polish journal of microbiology*, 65 (4), 7.
- Siddikee, M. A., Hamayun, M., Han, G. H., & Sa, T. M. (2010). Optimization of gibberellic acid production by *Methylobacterium oryzae* CBMB20. *Korean Journal of Soil Science and Fertilizer*, 43 (4), 522-527.
- Singh, N. B., Singh, A., Khare, S., Yadav, V., Bano, C., and Yadav, R. K. (2020). Mitigating Strategies of Gibberellins in Various Environmental Cues and Their Crosstalk with Other Hormonal Pathways in Plants: A Review. *Plant Molecular Biology Reporter*, 1-16.
- Sleem, D. A. E. (2013). Studies on the Bio production of Gibberellic Acid from Fungi. 7-120.
- Swetha, S., and Padmavathi, T. (2016). Study of acid phosphatase in solubilization of inorganic phosphates by *Piriformospora indica*. *Polish Journal of Microbiology*, 65 (4), 407-412.
- Ukale, D. U., Bhagwat, R. V., Upadhyay, S. K., Cukkemane, N., and Cukkemane, A. A. (2016). Metabolic analysis of liquid formulations of organic manures and its influence on growth and yield of *Solanum lycopersicum* L. (tomato) crop in field. *Biocatalysis and Agricultural Biotechnology*, 8, 50-54.
- Verma, P., Yadav, A. N., Kumar, V., Singh, D. P., and Saxena, A. K. (2017). Beneficial plant-microbes interactions: biodiversity of microbes from diverse extreme environments and its impact for crop improvement. *Plant-microbe interactions in agro-ecological perspectives* 543-580.

Estimation of Correlation and Path coefficient analysis for yield and its component traits in tomato

Vijay Kumar* and S. K. Singh

Dept. of Genetics and Plant Breeding (Ag Botany), C.C.R (PG) College, Muzaffarnagar, U.P. 251001

*Corresponding Author- vijayram9092@gmail.com

Abstract

An experiment was conducted at agriculture research farm, C.C.R. (PG) College Muzaffarnagar, U.P. to study the path and correlation which are very important association among yield and related component traits. Days to first flowering have significant genotypic and phenotypic correlation with number of primary branch (0.214, 0.189) and plant height (cm) (0.190, 0.150). Similarly, Days to first flowering, number of primary branches and number of fruits/plant show affirmative effect on fruit yield at genotypic and phenotypic levels. Further fruit yield can be improved by making the selection of desirable traits in important progenies of further crop improvement programmes.

Keyword: Tomato, Genotypic Correlation, Phenotypic Correlation and Path Analysis

INTRODUCTION:

Tomato (*Lycopersicon esculantum* Mill.) is a member of the family solanaceae and considered a valuable vegetable crop worldwide (He *et al.*, 2003). Vegetables are an important part of our diet as they provide energy and are also a rich source of minerals (Iron and phosphorus), vitamins (Vitamin A and C), organic acid, essential amino acids, dietary fibres, etc. Further tomato of fruit provides total sugar (3-4%), total solids (4-7%), ascorbic acid (15-30 mg/100g), Titratable acidity (7.5-10 mg/100ml) and fruit weight of lycopene (20-50mg/100g). Tomato and its products lower risk of developing digestive tract and prostate (Giovannucci *et al.*, 2002).

Path analysis and Correlation coefficient are key determinants to objective the association between the yield and related components. For the plant breeders positively correlated with the yield are important for selecting desirable genotype (Feyzian *et al.* 2009). All the changes in the components need not, however be expressed by changes in the yield. This may be due to degrees of correlations (positive and negative) among yield and its other components or own among the components themselves. In the present study correlation coefficients and path analysis relevance of direct and indirect association of all characters with the final yield and its components have been studied.

Material and Methods:

The study was conducted during year 2020-21 at Agriculture Research Farm, C.C.R. (PG) College Muzaffarnagar, U.P. The material for the present study comprised of ten parents of tomato, namely EC-620406, EC-163605, EC-

165700, EC-631364, EC-164563, EC-521067, EC-528360, EC-145057, Floradade, and Arka Meghali and forty five hybrids were produced through half diallel mating in Randomized Block Design with three replications. All the recommended package of practices was adopted for raising a healthy crop. Five plants were randomly selected from each treatment for observations. Analysis of covariance for all combinations was for to study correlations. Phenotypic/genotypic correlations were calculated by the formulae recommended by Al-Jibouri *et al.* (1958). Path analysis was calculated as per the procedure outlined by Wright (1921), and Dewey and Lu (1959). Phenotypic and genotypic correlations and path coefficient analysis were analyzed using R software (Version 4.1.1). Using Lavaan Package (Rosseel, 2012).

Result and Discussion:

In the present investigation the results revealed Table 1 and 2 that there is a significant and positive correlation of days to first flowering at genotypic level was observed with days to 50% flowering (0.951), no. of primary branches/plant (0.215), plant height (0.191) and no. of fruits/cluster (0.178). Days to 50% flowering exhibited significant positive genotypic correlation with no. of primary branches/plant (0.297), number of fruits/cluster (0.267), total yield/plant (0.195), no. of secondary branches/plant (0.174). No. of primary branches per plant at genotypic level was observed with no secondary branches (0.533), fruit diameter (0.436), days to 50% flowering (0.297) and days to first flowering (0.215). Plant height at genotypic level was observed with no. of secondary branches per plant, days to first

flowering, days to 50% flowering and pericarp thickness with values of 0.223, 0.159, 0.191, and 0.155, respectively. Average fruit weight at genotypic level was observed with total yield per plant (0.574), TSS (0.565), pericarp thickness (0.533), fruit diameter (0.513) and fruit length (0.482). Pericarp thickness at genotypic level was observed with fruit length (0.657), average fruit weight (0.533), TSS (0.451), fruit diameter (0.209), T.A (0.197) and plant height (0.156). Ascorbic acid at genotypic level was observed with days to 50% fruiting (0.293) and no. of fruits per cluster (0.151). Total yield per plant at genotypic level was observed with average fruit weight (0.574), no. of fruits per plants (0.462) TSS (0.413), T.A (0.340) and no. of fruits per cluster (0.259). The results are in line with the previous reports (Singh *et al.* 2018, Mishra *et al.* 2019, Kumar *et al.* 2020, Maurya *et al.* 2020 and Akhter *et al.* 2021).

A significant and positive correlation of days to first flowering at phenotypic level was observed with days to 50% flowering (0.849), number of primary branches/plant (0.190), number of fruits/cluster (0.154) and plant height (0.151). Days to 50% flowering exhibited significant positive phenotypic correlation with no. of primary branches/plant (0.194), no. of fruits/cluster (0.169) and days to first flowering (0.849). No. of primary branches/plant revealed significant positive phenotypic correlation with no. of secondary branches/plant (0.489), fruit diameter (mm) (0.305), days to first flowering (0.190), days to 50% flowering (0.194). Fruit

diameter showed significant positive genotypic correlation with average fruit weight (0.418), fruit length (0.290), TSS (0.181), total yield/plant (0.169), titratable acidity (0.168), pericarp thickness (0.165), number of primary branches/plant (0.305). Average fruit weight showed significant positive phenotypic correlation with total yield/plant (0.553), pericarp thickness (mm) (0.479), TSS (0.473), titratable acidity (0.214), fruit length (mm) (0.463) and fruit diameter (mm) (0.418). Total yield/plant (g) showed significant positive phenotypic correlation with average fruit weight (g) (0.553), no of fruit/plant (0.452), TSS (0.344), plant height (cm) (0.334), titratable acidity (0.328), number of fruits/cluster (0.193) and fruit diameter (0.169). The results are in line with the previous reports (Mishra *et al.* 2019 and Kumar *et al.* 2020).

The path coefficient is considered an important method for estimating the direct or indirect effect cause of inter/intra association. The similarly genotypic path coefficient analysis was also studied in Table 3 and Figure (1 & 2). The direct effect on total yield recorded to be the highest by average fruit weight (1.120) followed by no. of fruit/plant (1.006), days to first flowering (0.902), plant height (0.195), total yield/plant (0.164), fruit diameter (0.118), T.A. (0.095), pericarp thickness (0.074), ascorbic acid (0.054), no. of primary branches (0.045) and fruit length (0.044). While indirect effect at genotypic level days to first flowering had positive (indirect) effect on total yield via average




Figure 1. Path coefficient diagram and effect relationships among quantitative traits and total yield.

fruit weight (0.109), no. of fruit/plant (0.090), TSS (0.056), plant height (0.038), days to 50% fruiting (0.017), no. of primary branches (0.010) fruit diameter (0.007). Days to 50% fruiting had positive indirect effect on total yield via days to 50% flowering (0.194), TSS (0.060), plant height (0.023), no. of secondary branches (0.017), ascorbic acid (0.016), pericarp thickness (0.013). No. of secondary branches per plant had indirect effect on total yield via no. of fruit/plant (0.172), days to first flowering (0.116), days to 50% fruiting (0.058), plant height (0.044), TSS (0.033), no. of primary branches (0.024) and Fruit diameter (0.018). Fruit length had positive indirect effect on fruit yield via average fruit weight (0.539), days to 50% flowering (0.176), pericarp thickness (0.049), fruit diameter (0.041), T.A. (0.032), days to 50% fruiting (0.015), no. of secondary branches (0.009) and fruit/cluster (0.007). Pericarp thickness (mm) had positive (indirect) effect on total yield via average fruit weight (0.597), plant height (0.031), fruit length (0.029), fruit diameter (0.025), T.A. (0.019), no. of primary branches per plant (0.014), ascorbic acid (0.003). Ascorbic acid had positive (indirect) effect on total yield via days to 50% flowering (0.148), TSS (0.132), no. of fruit/plant (0.035), no. of secondary branches per plant (0.006). The finding of these characters is supported by Jogi *et al.* (2018), Namdev and Dongre (2019), Kumar *et al.* (2020) and Singh *et al.* (2021).

Phenotypic path analysis direct and indirect effect on different traits present in table 4. The direct effect on total yield recorded to be highest by average fruit weight (0.800) followed by no. of fruit/plant (0.757), ascorbic acid (0.245), total yield/plant (0.137), TSS (0.085), no. of primary branches (0.054), days to first flowering (0.039), T.A. (0.038), plant height (0.022) and fruit length (0.021). While (indirect) effect at phenotypic level days to first flowering had positive (indirect) effect on total yield via average fruit weight (0.069), number of fruit/plant (0.063) and days to 50% fruiting (0.010). Days to 50% flowering had positive (indirect) effect on total yield via no. of fruit/plant (0.088), average fruit weight (0.073), days to first flowering (0.033), days to 50% fruiting (0.015), no. of primary branches (0.011) and plant height (0.003) show positive (indirect) effect on total yield recorded. Similarly, plant height (cm) also had positive (indirect) effect on total yield via days to first flowering (0.007), no. of fruit/cluster (0.005) and fruit diameter (0.003). Fruit diameter (mm) had positive (indirect) effect on total yield per plant via average fruit weight (0.335), TSS (0.016), no. of primary branches (0.017), days to 50% fruiting (0.009), T.A. (0.007), fruit length (0.006), number of fruit/cluster (0.003) and first flowering (0.002) show positive (indirect) effect on total yield recorded. Pericarp thickness (mm) show positive (indirect) effect on total yield via average fruit weight

Figure 2: Principal Component Analysis (PCA) Biplot of tomato.

