

Nitrogen and potassium requirements of processing variety of potato Kufri chipsona 3 under North Gujarat condition

H. L. CHAUDHARI¹, P. P. CHAUDHARI² AND C. K. PATEL³

Department of Agronomy, C. P. College of Agriculture,
S. D. Agricultural University, Sardarkrushinagar - 385 506 (Gujarat)

Received : 03 June, 2015

ABSTRACT

The field experiment was conducted during *rabi* season of the year 2012-13 at Potato Research Station, Sardarkrushinagar Dantiwada Agricultural University, Deesa with nine treatment combinations consisting of three levels of nitrogen viz., 225 kg N/ha, 275 kg N/ha and 325 kg N/ha and three levels of potassium viz., 225 kg K₂O/ha, 275 kg K₂O/ha and 325 kg K₂O/ha tested under factorial randomized block design with four replications. The yield of 'A' and 'B' grade tubers was increased upto 275 kg N/ha. nitrogen levels brought significant variation in chips colour index and content of reducing sugar. In case of Potassium, Potassium levels 325 kg K₂O/ha and 275 kg K₂O/ha behaved equally and produced significantly higher weight of 'A' and 'B' grade tuber with minimum weight of 'C' grade tubers. Maximum reducing sugar content and chips color index score were observed with the application of 325 kg K₂O/ha however increase was upto 275 kg K₂O/ha. Thus, for getting higher production of processing grade tuber yield and net realization, potato cv., Kufri Chipsona 3 should be fertilized with 275 kg N/ha and 275 kg K₂O/ha.

Key words : Potato, processing, nitrogen, potassium, tuber yield, chips colour

Increasing trends in potato production cannot be sustained without diversification in its utilization. Potato processing industry therefore has to be essential to sustain present growth rate. Potato grown in cooler North-Western plains have lower dry matter and high reducing sugar reflecting products, hence unfit for processing. In Gujarat warm dry and mild night temperature during the crop season favours for production of high dry matter and accumulation of low sugars. Therefore, the region is ideal for potato processing. This could help in reduction of market glut at peak season of harvest. Thus, commercialization of agriculture and stress on value addition through processing, have led to evaluate the newly released processing varieties for their chips quality and yield under Gujarat condition.

Hence a study was undertaken to standardize the nitrogen and potassium requirements of chipping cv

Kufri Chipsona 3 of potato for getting higher processing grade tubers yield and quality.

MATERIALS AND METHODS

The field experiment was conducted during *rabi* season of the year 2012-13 at Potato Research Station, Sardarkrushinagar Dantiwada Agricultural University, Deesa (24° 12' North latitude and 72° 12' East longitude with an elevation of 136.4 meters above the mean sea level). Nine treatment combinations consisting of three levels of nitrogen viz., N₁: 225 kg N/ha, N₂: 275 kg N/ha and N₃: 325 kg N/ha and three levels of potassium viz., K₁: 225 kg K₂O/ha, K₂: 275 kg K₂O/ha and K₃: 325 kg K₂O/ha were tested under factorial randomized block design with four replications. The soils of the experimental field was loamy sand in texture having good drainage, low in available nitrogen (192.0 kg/ha), medium in available phosphorus (26.7 kg/ha) and high in potash (346.0 kg/ha). The experimental crop cv Kufri Chipsona-3 was planted on 29th November. Potato tubers of variety Kufri

¹P.G. Student, ²Associate Professor, ³Associate Professor,
College of Agriculture, NAU, Bharuch

Chipsona 3 were cut in pieces, keeping two to three eye bud with approximately 25 to 40 g weight and planted at the spacing of 50 X 20 cm spacing. Half dose of N and full dose K (As per treatments) along with full recommended dose of phosphorus (140 kg P_2O_5 /ha) were placed in bends below the seed pieces at the time of planting and remaining half dose of N was applied at earthing up 30 days after planting (DAP). Total eleven irrigations (approximately 50 mm each) were given to the experimental plot. The irrigations were discontinued one week before haulm killing. The dehaulming was done mechanically at full maturity and harvesting was done two weeks later. Tubers were graded in to three categories viz., 'A' grade (Large size, > 60 g), 'B' grade (Medium size, 20-60 g) and 'C' grade (Small size, <20 g). For determining chip colour ten tubers from each plot were washed, hand peeled and cut into 1.75 mm thick slices. Slices were washed in normal water to remove surface starch and then the slices were dried on paper towels. The dried slices were fried in edible oil at 180°C in a deep fat fryer (Moulinex make) till the bubbling stopped. The fried chips were drained to remove excess oil and then graded from 1 to 10 (1 being lightest and 10 the darkest, the chip colour grad up to 3 is considered acceptable) under the fluorescent tube light using chip colour cards (Ezekiel *et al.*, 2003). The specific gravity of potato was calculated as:

$$\text{Specific gravity of potato (g/cc)} = \frac{\text{Specific gravity of water} \times 5000}{5000 - \text{WIW}}$$

Five tubers from each plot were chopped and mixed and prepared composite mixed samples and it put into the boiling isopropyl alcohol (80 per cent v/v) in a flat bottomed flask. The samples were than boiled at 70°C for two minutes for fixing. The fixed samples were put to reflux at 70°C twice, and filtrate through Watman No. 1 filter paper. The filtrate was evaporated at 70°C to concentrate to 10 ml. The concentrated extract was diluted to 100 ml with double distilled water to make the final extract. The extract was used to determine the reducing sugar content by Nelson's method(1944).

RESULTS AND DISCUSSION

Weight of tubers grade viz., A, B and C were influenced significantly due to different nitrogen levels (Table 1). Both the higher levels of nitrogen N_2 (275 kg N/ha) and N_3 (325 kg N/ha) performed equally and

significantly the lowest value for 'A' and 'B' grade tuber yield were observed with N_1 , whereas, the lowest values of 'C' grade tuber yield were beared by N_3 (325 kg N/ha)(Table 1). This indicate that curtailing of dose of up to 25 per cent optimized the nitrogen requirement to produced quality potato. This might be due to better assimilation of carbohydrate to the tubers, which ultimately helped in the enlargement of the tuber size and weight of tuber as reported by Malik *et al.*, (2002).

Nitrogen application of N_2 (275 kg N/ha) and N_3 (325 kg N/ha) realized spectacular improvement in total tubers yield of potato and increased total tubers yield to the tune of 12.99 and 13.02 per cent over N_1 (225 kg N/ha)(Table 1). Higher rate of nitrogen application provides better growth, development and translocation of photosynthesis from source to sink (tuber), which resulted in to higher yield of large size tubers, concluded by Patel and Patel (2001). These finding corroborate the reports of Singh *et al.* (2002) and Vijaya Lakshmi, *et al.* (2011). Nitrogen application @ 275 kg N/ha recorded highest gross and net realization of ₹ 289035 /ha.

Conspicuous difference in weight of different grades of tuber were observed due to potassium application (Table 1). Potassium levels K_3 (325 kg K_2O /ha) and K_2 (275 kg K_2O /ha) performed equally good and produced significantly higher weight of 'A' and 'B' grade tubers with minimum weight of 'C' grade tubers. The lowest value for 'A' and 'B' grade tuber yield were observed with K_1 (225 kg K_2O /ha) whereas, for 'C' grade tuber yield the lowest values were beared by K_3 (325 kg K_2O /ha). Potassium application @ 275 kg K_2O /ha recorded highest net realization of ₹ 279593 /ha. These finding corroborate the reports of Parveen Kumar *et al.*, (2004). Potassium promote large size tubers by increasing water accumulation in tubers (Vijya Lakshmi *et al.*, 2011).

Among the potassium levels, K_3 (325 kg K_2O /ha) and K_2 (275 kg K_2O /ha) did not differ from each other, and produced 8.61 and 8.37 per cent higher total tubers yield over K_1 (225 kg K_2O /ha) respectively (Table 1). Higher rate of potassium provides better yield attributes and translocation of photosynthesis from source to sink (tuber) which resulted in to higher yield of large size tubers. These finding corroborate the reports of Khandakhar *et al.*, (2004).

Nitrogen application did not show positive response on specific gravity of potato (Table 1). However, higher

Table 1 Grade wise total tuber yield (q/ha), net realization and quality parameters of potato as influenced by nitrogen and potassium fertilization

Treatments	Tuber yield (q/ha)				Net realization (₹ /ha)	Chips colour index (0-10 grade)	Specific gravity (gm/cc)	Reducing sugar mg/100 gm fresh wt.
	A grade	B grade	C grade	Total				
Nitrogen (Kg/ha)								
N ₁ : 225	124.06	115.30	59.74	301.85	248471	1.42	1.059	55.48
N ₂ : 275	142.14	145.38	50.74	341.09	289036	1.56	1.051	62.22
N ₃ : 325	143.29	148.61	46.48	341.17	288481	1.68	1.058	66.21
S. Em. ±	3.95	3.70	1.64	7.65		0.04	0.031	1.99
C.D. at 5%	11.53	10.80	4.78	22.32		0.13	NS	5.81
Potassium(Kg/ha)								
K ₁ : 225	126.21	126.63	54.95	310.46	253763	1.44	1.059	56.59
K ₂ : 275	140.63	139.02	53.90	336.45	279593	1.59	1.045	62.93
K ₃ : 325	142.65	143.63	48.11	337.20	278913	1.63	1.064	64.40
S. Em. ±	3.95	3.70	1.64	7.65		0.04	0.031	1.99
C.D. at 5%	11.53	10.80	4.78	22.32		0.13	NS	5.81
Interaction								
N × K	NS	NS	NS	NS		Sig	NS	NS
C.V. %	10.03	9.39	10.84	8.08		9.64	10.27	11.24

Table 2 Interaction effect of nitrogen and potassium on chips colour index potato

Nitrogen (Kg/ha)	Potassium (Kg/ha)		
	K ₁ :225	K ₂ :275	K ₃ :325
N ₁ :225	1.38	1.38	1.51
N ₂ :275	1.45	1.72	1.50
N ₃ :325	1.49	1.67	1.88
S. Em. ±	0.08		
C.D.	0.20		
C.V %	9.64		

levels of nitrogen realized spectacular improvement in reducing sugar content (4.10). Maximum content of reducing sugar was noted with N₃ (325 kg N/ha), which accounted 19.34 and 6.41 per cent higher reducing sugar content than N₂ (275 kg N/ha) and N₁ (225 kg N/ha), respectively. The score of chips colour index increased significantly as the rate of application of nitrogen increase up to N₂ (275 kg N/ha). Nitrogen N₃ (325 kg N/ha) had higher value of chips colour index, which means more darkness in chips, whereas level N₁ (225 kg N/ha) had lower colour index. This finding is well supported by Swiniarski and Ladenberger (1970).

From the data (Table 1), it appeared that reducing sugar content was linearly increased with increase in potassium level. Potassium level K₃ (325 kg K₂O/ha) and K₂ (275 kg K₂O/ha) had produced tuber having 13.80 and 11.20 per cent higher reducing sugar content than that of produced by K₁ (225 kg K₂O/ha), respectively. These finding corroborate the reports of Ackarya and Mondal (2007). Potassium levels brought significant difference in chips colour index (Table 2). The maximum and minimum score values of chips colour index were noted with K₃ (325 kg K₂O/ha) and K₁ (225 kg K₂O/ha), respectively. Potassium levels K₃ and K₂ did not differ each other and were significantly higher than K₁. These finding are well supported by Khan *et al.*, (2012).

Interaction between nitrogen and potassium levels (N×K) was found to be significant in respect of chips colour index of potato (Table 2). Chips colour index score increased with increase in N levels with correspondence increase in K₂O level. Treatment combination N₃K₃ (325 kg N/ha with 325 kg K₂O/ha) recorded significantly the highest value of chips colour index (1.88) followed by treatment N₂K₂ (1.72). These results are in accordance with those reported by Ishwori *et al.*, (2011).

CONCLUSION

Based on the results from one year experimentation it is concluded that to get higher production of processing grade tuber yield with maximum chips color index and reducing sugar content as well as net realization, potato cv., Kufri Chipsona-3 should be fertilized with 275 kg N/ha and 275 kg K₂O/ha.

LITERATURE CITED

- Ackarya, D and S S, Mondal 2007. Effect of integrated nutrient management on the potassium content in the plant and its effect on the quality characters and disease infestation of potato at kalyani, West Bengal. *Indian Journal of Agricultural Sciences*, **77**(10): 2007
- Ezekiel, R., Singh, B. and Kumad, D. 2003 A reference chart for potato chip colour for use in India. *Journal of Indian Potato Association*, **30**: 259-265
- Ishwori, P., Gautam, Bhim, B. Khatri, Duryodhan, Chaudhary, 2011. Effect of nitrogen and potassium on yield, storability and post harvest processing qualities of potato for chips. *Nepal Agricultural Research Journal*, **11**:40-51
- Khandakhar, S.M.A.T., Rahman, MM., Uddin, M.J., Khan, S.A., Quddus, K.G. 2004. Effect of lime and potassium on potato yield in acid soil *Pakistan Journal of Biological Sciences*, **7**: 380-383
- Khan, M., Ehsanakhtar, M. M., Masud Mahmood, M., Naeemsafdar, M., 2012. Potato tuber yield and quality as affected by rates and sources of potassium fertilizer. *Journal of Plant Nutrition*, **35**: 664-677
- Malik, Y.S., Bhatiya, A.K., Singh, N., Nehra, B.K. and Khurana, S.C. 2002. Effect of nitrogen, seed size and spacing on seed potato production in cv. Kufri Suttlej. In, Potato, Global Research and development, (Proceedings of Global Conference on Potato, New Delhi, 6-11 December 1999) Khurana SMP, Shekhawat GS, Pandey SK and Singh BP (Eds.) II:861-865.
- Nelson, N. 1944. A photometric adoption of the somogyi method for the determination of glucose. *Journal of Biological Chemistry*, **153**: 375-80.
- Parveen, Kumar, Pandey, S.K., Singh, S.V., Sanjay, Rawal, Kumar, D. 2004. Effect of potassium fertilization on processing grade tuber yield and quality parameters in potato. *Indian Journal of Agricultural Science*, **74** (4):177-179
- Patel, J.C. and Patel, B.K. (2001) Response of potato to nitrogen under drip and furrow methods of irrigation. *Journal of Indian Potato Association*, **28** (2/4): 293-295.
- Singh, N., Sood, M.C. and Sharma, R.C. 2002. Effect of irrigation levels, cultural practices and nitrogen application on potato production under drip and sprinkler method of irrigation. Potato global research and development-proceeding of the Global Conference on potato. New Delhi, 6-11, December, 923-925
- Swinorski, E. and Ladenberger, D. 1970. The sugar content of potato tubers grown with different rates of nitrogen application. *Potato Research*, **13**: 114-118.
- Vijaya Lakshmi, D., Padmaja, G. and Chandrasekhar, P. Rao, 2011. Effect of levels of nitrogen and potassium on yield, uptake and economics of potato grown on alfisols of Andhra Pradesh. *The Journal of Research ANGRAU*, **39**(4) 59-62.