

(0.384), TSS (0.028), fruit length (0.013), T.A. (0.010), ascorbic acid (0.008), plant length (0.004) and days to first flowering (0.002) show positive (indirect) effect on total yield recorded. Total Soluble Solid had positive (indirect) effect on total yield recorded by average fruit weight (0.379), titratable acidity (0.024), days to 50% fruiting (0.012), no. of fruits/cluster (0.009), no. of secondary branches (0.005) and days to 50% flowering (0.001) positive (indirect) effect on total yield. Ascorbic acid show positive (indirect) effect on total yield via number of fruit/plant (0.023), no. of secondary branches (0.003), days to 50% flowering (0.002). Titratable acidity show positive (indirect) effect on total yield via no. of fruit/plant (0.173), average fruit weight (0.170), TSS (0.052), days to 50% fruiting (0.009), fruit length (0.007), no. of secondary branches (0.002), no. of fruit/cluster (0.001) and days to 50% flowering (0.001), which fall in line with same previous results (Namdev and Dongre 2019, Kumar *et al.* 2020 and Akhter *et al.* 2021).

From present investigation it may be concluded that, no. of fruit / cluster, no. of fruits/plant, and fruit diameter and weight were the major yield contributing characters which showed positive and significant association with total yield /plant and also exhibited high direct effect on total yield /plant. Therefore, due there is a need to study characters in for selection which would help in isolating high yielding genotypes from highly segregating population to improve yield potential of tomato.

REFERENCES

- Akhter, M., Fariha, N. A., Muhammad, M., Rashid, B. and Naheed, Z. 2021. Genetic Variability, Correlation Coefficient, Path Coefficient and Principal Component Analysis in Tomato (*Solanum lycopersicum* L.). *Plant Cell Biotechnology and Molecular Biology* 22:46-59.
- Al-Jibouri, H. A., Miller, P. A. and Robinson, H, F. 1958. Genotypic and environmental variances and co-variances in an upland cotton cross of interspecific origin. *Agronomy Journal*, 50: 633-636.
- Dewey, D. R. and Lu, K. H. 1959. A correlation and path coefficient analysis of components of crested wheat grass seed production, *Agronomy Journal*, 51(9):515-518.
- Feyzian, E., Dehghani, H., Rezai, A. M. and Javaran M. J. 2009. Diallel cross analysis for maturity and yield-related traits in melon (*Cucumis melo* L.). *Euphytica*, 168: 215-22.
- Giovannucci, E., Rimm, E.B., Liu, Y., Stamper M.J. and Willett, W.C. 2002. A prospective study of tomato products, lycopene, and prostate cancer risk. *J. National Cancer Institute*. 94 (5): 391-398.
- He, C., Poysa, V. and Yu, K. 2003. Development and characterization of simple sequence repeat (SSR) markers and their use in determining relationships among *Lycopersicon esculentum* cultivars. *Theoretical Applied Genetics*, 106: 363-373.
- Jogi, M., Lingaiah, H. B., Indires, K. M., Singh, T. H., Samuel, D. K. and Ramachandra, R. K. 2018. Genetic divergences studies in tomato. *International Journal of Current Micro Applied Science* 7: 2231-2237.
- Kumar J, Singh Nirajan, Singh Pooshpendra, Yadav Lalit, Singh Budhesh Pratap and Tomar S. 2020. Correlation and path analysis in tomato (*Solanum lycopersicon* L.). *International Journal of Chemical Studies* 9 (1):3158-3160.
- Maurya, V. K., Kumar, S., Prasad, A. K., Bhatt, M. L. B. and Saxena, S. K. 2020. Structure-based drug designing for potential antiviral activity of selected natural products from Ayurveda against SARS-CoV-2 spike glycoprotein and its cellular receptor. *Virus Disease* 31:179-193.
- Namdev, S. K., and Dongre, R. 2019. Correlation and Path Analysis in Tomato. *Research Journal of Agricultural Sciences* 9:588-590.
- Wright, S. 1921. Correlation and causation. *Journal of Agriculture Resources* 66(4); 557-585.
- Yves Roosseel. 2012. Lavaan: An R package for structural equation modeling. *Journal of Statistical Software* 48 (2): 1-36.

Study on horticultural traits and incidence of physiological disorders of indigenous seedling mango germplasm in central and sub-montane zones of Punjab

Gurbir Singh¹ and Sukhdev Singh²

¹Faculty of Agriculture and Forestry, Guru Nanak Dev University Amritsar, Punjab, India.

² Director of Research, Khalsa College, Amritsar, Guru Nanak Dev University Amritsar, Punjab, India.

Corresponding author: gurbirsingh011@gmail.com

Abstract

A Survey on horticultural traits and incidence of physiological disorders of seedling mango germplasm was carried out in central and sub-montane zones of Punjab, during February to October 2012 (on year) and February to October 2013 (off year). The superior genotypes of seedling mangoes to peculiar character were selected on the basis of information provided by tree owners, local inhabitants and officials from Department of Horticulture, Punjab. Wide variability was recorded for eating quality in fruits at the time of ripening indicates that the fruit storage life in different evaluated seedling mango germplasm varied from 3 to 5 days. The time of fruit maturity also differed amongst the genetic resources under evaluation. The seedling mango germplasm collected in central and sub-montane zone of Punjab were irregular, alternate and regular bearers. A wide range of variability was observed in fruit yield of collected germplasm. Thus, there is tremendous possibility of such material to act as source for further propagation as well as breeding material for hybridization to evolve new varieties with desirable traits.

Key words: *Mangifera indica*, fruit quality, yield, variability, Punjab; India introduction

INTRODUCTION:

Mango ! belongs to genus *Mangifera* under family Anacardiaceae and order Sapindales comprising of 69 species (Kostermans and Bompard, 1993), which are distributed throughout the world. Mango peel and pulp contain carotenoid pigments, vitamin C, polyphenols, omega-3 poly unsaturated fatty acids, proteins and carbohydrates along with minerals like calcium, phosphorus and iron. Fruit is utilized at all stages of its development. Young and unripe fruits are liked because of their acidic taste and are used for culinary purposes as well as for pickle, chutney and amchoor (mango powder) making. Ripe fruits are utilized in preparing squash, jam, jelly, custard powder, mango leather (Amb papad) and mango toffee. Its wood is used for making furniture, packing

boxes, match boxes and also as a source of fuel. It is the national fruit of India, Pakistan and the Philippines and the national tree of Bangladesh. The heavy canopy of the seedling mango is a source of shelter and shade for both animals and humans and also good as a wind break against strong winds. No other fruit can be put to so many diversified uses in the form of processed product as the mango. Therefore, it is rightly acknowledged as the king of fruits in India. It enjoys the same popularity in the tropics as apple in the temperate regions.

Mango is grown commercially in India, Indonesia, Bangladesh, Philippines, Malaysia, Sri Lanka, Thailand, Pakistan, Egypt, Sudan, South Africa, West Indies, Hawaii, Brazil and USA (Florida) etc. In India it is successfully growing in the states of Uttar Pradesh, Andhra Pradesh,

Bihar, West Bengal, Kerala, Karnataka, Punjab and Haryana. The mango is now cultivated mostly in frost-free tropical and warm subtropical climates, almost half of the world's mangoes are cultivated in India alone.

India is the largest producer of choicest table varieties of mango in the world and wide variability has been found in fruit shape, size and taste but commercial importance is given to nearly about 30 mango cultivars in the country. But still, mango industry lags far behind in the world due to low productivity, higher incidence of insect pests, diseases and physiological disorders etc. Mango is cross pollinated fruit crop and almost majority of mango cultivars under cultivation in the world are selections from open pollinated seedling populations.

MATERIAL AND METHOD The present study was conducted in central and sub-montane zones of Punjab consecutively for three years during February to October 2012 (on year) and February to October 2013 (off year) and February to October 2014 (on year). More than hundred genotypes were selected on the basis of information provided by tree owners, local inhabitants and officials from Department of Horticulture, Punjab. Out of which twenty five superior genotype of seedling Mangoes to peculiar character were selected. The selected genotypes were evaluated with the help of IPGRI, Rome 'Mango Descriptor' for various vegetative, floral, fruit maturity, disorders and storage characters etc. Thirty fruits/sample were randomly selected from all the sides of the tree

Yield/tree: The yield/tree was recorded at optimum maturity in kg for estimating yield of a particular tree, mango fruits were harvested and weighed on a physical balance and registered according to their providing marking.

Fruit maturity period: The time of maturity was dependent upon various external characters like appearance of waxy coating, dots on the fruits and relative size of the fruit, change in colour and firmness. the natural fall of mature fruit is

considered to be the indication of fruit maturity in mango. All the fruit samples for physico-chemical analysis were taken at optimum maturity. The maturity period for the evaluated types was recorded and classified as : (a) Early (b) Medium (c) Late

Fruit storage life: It was recorded number of days of storage of ripe fruits under ambient conditions after harvest.

Eating quality: The combined assessment of flavour/acidity, sweetness, aroma and astringency of TSS, acidity and sugars to be made and classified as : (a) Poor (b) Good (c) Very good (d) Excellent

Bearing behaviour: The regularity of bearing was assessed during the two successive years of the research study. However the knowledge about regularity of bearing in the past was also taken from the owner and employees of the orchards and are classified as: (a) Irregular (b) Regular (c) Alternate

Horticultural traits and incidence of physiological disorders

Results with respect to various horticultural traits such as eating quality, fruit storage life, time of fruit maturity, bearing behaviour and yield per tree are documented in Table – 1. The genotypes AA-7, AA-9, AA-42 and ABG-1 had excellent taste, flavour and aroma. Genotypes AA-4, AA-16, AA-29 and AA-39 had very good and rest of the genotypes having good taste, flavour and aroma. These results lend support from findings of earlier research workers Iyer and Subramanyam (1993) and Desai *et al.* (2000). The highest keeping quality was recorded in genotypes AA-3, AA-8, AA-39 and AR-1 in which fruit remained sound condition for 5 days. Variation in fruit maturity in different seedling mango germplasm might be due to inherent genetic variation. The genotypes AA-1, AA-2, AA-3, AA-8 and AA-19 mature during 1st week of July and considered as early maturing genotypes. Late maturing genotypes which matured during 3rd and 4th week

of July were AA-7, AA-9, AA-39, AA-42, ABG-1, AGKB-64, AP-1, AUG-1 and AR-1. These findings are corroborated well with experimental outcomes of Boshra *et al.* (2007), Sharma *et al.* (2007) and Guirado *et al.* (2009). Most of the seedling mango germplasm evaluated in the central and sub-mountain zones of Punjab were found to be irregular and/or alternative bearer in nature. However, genotypes AA-101, ABG-1 and AGKB-64 showed tendency towards regular bearing. The results regarding bearing habit are in good tune with the research findings of Saha (1972) and Singh (1988). The genotype AKS-1 ranked first with yield (1400.20 kg/tree), followed by AA-39 (497.84 kg/tree), AA-29 (397.75 kg/tree) AA-19 (385.96 kg/tree), respectively, were high yielding genotypes. The minimum yield (79.50 kg/tree) was found in genotype GRB-1. The findings obtained in the present study lends support from the previous reports of Sanjay and Singh (2004) and Rao and Rao (2007), who also found high range of variation in yield per tree. The variation in the yield of evaluated seedling mango genotypes in the current investigation might be attributed to difference in growing conditions, genetic variations and cultural practices that ultimately influenced the size of the plant.

Fruits from evaluated germplasm were tested for incidence of disorders like malformation, black tip and spongy tissue. Considerable variation had been noticed with respect to occurrence of mango malformation; however none of the genotypes apparently exhibited the incidence of black tip and spongy tissue (Table – 2). The least occurrence of malformation (4.10%) was observed in genotype AR-1, it was followed by the genotypes AUG-1 (5.3%). Genotypes AP-1, AA-16, AKS-1 and AA-39 were also found comparatively less prone to mango malformation and considered as lightly affected genotypes. The variation in genotype susceptibility to mango malformation might be due to genetic makeup, season, climate and tree age. This result of the present study is in good confirmation with the earlier findings of Dhar

and Gupta (1979), who recorded the incidence of malformation as low as 3.2% in mango under U.P condition. High range of variations in the occurrence of mango malformation also had been documented from investigations of Singh and Mandal (1999), Pinto *et al.* (2000) and Prakash *et al.* (2006).

CONCLUSION

From the overall outcomes of the present investigation it is evident that the germplasm with desirable traits can be proved to be good genetic material for the utilization in future breeding programmes for improving the mango varieties in Punjab. Genotypes AKS-1 along with AA-39, AA-29 and AA-19 were considered as high yielding genotypes. Amongst the evaluated germplasm AA-101, ABG-1 and AGKB-64 showed tendency towards regular bearing, where as genotypes AR-1 and AUG-1 showed the maximum resistance against mango malformation. Hence, genotypes such as AKS-1, GJB-1, GQ-1, AA-19, AA-29, AA-39, AA-42, AA-101, ABG-1, AGKB-64, AR-1 and AUG-1 can be included for future systematic breeding as well as hybridization programme of mango to inculcate and/or concentrate favourable attributes among the improved progenies.

REFERENCE

- Boshra, S.E., Abul-Soad, A.A. and Said, A.G. (2007). Selection of some mango seedlings grown in Assiut. *Egyptian Journal of Plant Breeding*, 11(1): 223-234.
- Desai, A.R. and Dhandor, D.G. (2000). Variation in Physico-chemical and morphogenetic characters of some mango varieties of Goa. *Acta Horticulture*, 509: 243-252.
- Dhar, L., Gupta, J.H. and Ram, P.B. (1979). A note on the performance of mango cultivars against malformation in Uttar Pradesh. *Prog. Hort.*, 11(3): 45-47.
- Guirado, E., Farre, J.M. and Nermoso, J.M (2009). Mango cultivar studies in Southern Mainland Spain. *Acta Hort.*, 820: 127-132.

- Iyer, C.P.A. and Subramanyam, M.D. (1993). Improvement of mango. *Advances in Hort. (Eds: KL Chadha and O.P. Pareek)*, 1: 267-278.
- Parkash, Satya, Sirohi, S.C. Rana, P. and Singh, R. (2006). Response of malformation to mango cultivars. *J. Res., SKUAST*. 5(1): 115-117.
- Pinto, A.C., Ramos, V.H.V. and Junqueira, N.T.V. (2000). New varieties and hybrid selections from mango hybridization program in central region of Brazil. *Acta Hort.*, 509: 207-212.
- Rao, K.D. and Rao, D.S.K. (2007). Evaluation of mango varieties and hybrids performance in lateritic soils under rainfed conditions. *Agri. Sci. Digest*, 27(4): 297-298.
- Saha, A.K. (1972) studies on some commercial characters of important early mango of Marshidabad district of West Bengal, India. *Acta Hort.*, 24: 39-54.
- Sanjay and Singh, J. (2004). Evaluation of mango hybrids for flowering, fruiting and fruit quality attributes. *Prog. Hort.*, 36(2): 343-346.
- Sharma, S.K. and Vishal, R. (2007). Studies on harvest maturity of 'Dashehari' and 'Langra' cultivars of mango growing under the sub-mountain region of Himachal Pradesh, India, *Asian J. Hort.*, 2(2): 37-42.
- Singh, J.P. and Mandal, B.K. (1999). Floral malformation varieties in Vaishali district of Bihar. *J. of Applied Biology*, 9(2): 159-162.
- Singh, R. (1988). Fruits. *National Book Trust*, New Delhi, 18-40.

Table 1: Horticultural traits of evaluated genotypes.

Selection number	Collector code	Eating quality	Fruit storage life (Days)	Time of fruit maturity	Bearing behaviour	Yield/ tree (kg)
1	AP- 1	Good	3.00	4th week of July	Irregular	158.20
2	AKCA - 14	Good	5.00	2nd week of July	Alternate	149.30
3	GJB -1	Good	3.00	2nd week of July	Irregular	189.10
4	AA -101	Good	4.00	2nd week of July	Regular	179.20
5	AA-15	Good	4.00	2nd week of July	Irregular	297.10
6	AA-4	Very Good	4.00	2nd week of July	Irregular	349.50
7	AA -16	Very Good	3.00	2nd week of July	Irregular	291.40
8	AA -42	Excellent	5.00	4th week of July	Irregular	247.30
9	GRB-1	Good	3.00	2nd week of July	Irregular	79.50
10	AA-3	Good	5.00	1st week of July	Irregular	345.15
11	GQ-3	Good	4.00	2nd week of July	Irregular	179.20
12	AA-20	Good	4.00	2nd week of July	Irregular	296.10
13	AUG.1	Good	3.00	4th week of July	Alternate	195.20
14	AA-9	Excellent	4.00	3rd week of July	Irregular	238.72
15	ABG-1	Excellent	4.00	3rd week of July	Regular	168.40
16	AGKB-64	Good	4.00	4th week of July	Regular	169.10
17	AA-29	Very Good	3.00	2nd week of July	Irregular	397.75
18	AA-39	Very Good	5.00	3rd week of July	Irregular	497.84
19	AA-2	Good	4.00	1st week of July	Irregular	349.41
20	AR-1	Good	5.00	4th week of July	Irregular	128.20
21	AA-8	Good	5.00	1st week of July	Irregular	287.76